

## APC Anti-Human CD122(TU27)

### Product Information

|              |  |
|--------------|--|
| Catalog No.: | APC-30051  |
| Conjugation: | APC  |
| Size:        | 50T/100T   |
| Form:        | liquid   |
| Host:        | Mouse  |
| Clonality:   | monoclonal   |
| Clone ID:    | TU27   |
| IsoType:     | IgG1, kappa  |
| Storage:     | PBS with 0.1% sodium azide, 1%BSA, pH 7.3, 2-8°C for 12 months<br>(Avoid repeated freeze / thaw cycles.) |

### Immunogen information


|              |  |
|--------------|--|
| Synonyms:    | Interleukin-2 receptor subunit beta IL-2RB p75 CD122 IL-2R $\beta$ |
| Uniprot ID : | P14784   |

### Application

|                     |       |
|---------------------|-------|
| Reactivity:         | Human |
| Tested Application: | FC    |

Recommended dilution: Volume per test: 5 $\mu$ L. Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Image:



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD56 Antibody and APC Anti-Human CD122 Antibody(Left). Lymphocytes are stained with FITC Anti-Human CD56 Antibody and APC Mouse IgG1,  $\kappa$  Isotype Control (Right).

