

## FITC Anti-Mouse CD16/32 Antibody(2.4G2)

### Product Information

|              |  |
|--------------|--|
| Catalog No.: | FITC-30106   |
| Conjugation: | FITC   |
| Size:        | 50T/100T   |
| Form:        | liquid   |
| Host:        | Rat  |
| Clonality:   | monoclonal   |
| Clone ID:    | 2.4G2  |
| IsoType:     | IgG2b, kappa   |
| Storage:     | PBS with 0.1% sodium azide, 1%BSA, pH 7.3, 2-8°C for 12 months<br>(Avoid repeated freeze / thaw cycles.) |

### Immunogen information


|              |   |
|--------------|---|
| Synonyms:    | CD16a/b CD32 CD32A/B FCG2A FCGR2A/B FCGR3 FCGR3A/B Fc fragment of IgG low affinity IIIa/b receptor Fc fragment of IgG low affinity IIIb receptor Fc fragment of IgG low affinity IIa/b receptor Fc gamma RIIa/b  Fc gamma receptor III A/B FcGR CD16/32 |
| Uniprot ID : | P08508 P08101   |

### Application

|                     |       |
|---------------------|-------|
| Reactivity:         | Mouse |
| Tested Application: | FC    |

Recommended dilution: Volume per test: 5µL. Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Image:



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD16/32 Antibody (filled red histogram). Unstained splenocytes (empty black histogram) are used as control.