

CBL antibody

Product Information

Catalog No.:	FNab01317
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5' UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause of Noonan syndrome-like disorder.

Immunogen information

Immunogen:	Cas-Br-M (murine) ecotropic retroviral transforming sequence
Synonyms:	E3 ubiquitin-protein ligase CBL Casitas B-lineage lymphoma proto-oncogene Proto-oncogene c-Cbl RING finger protein 55 RING-type E3 ubiquitin transferase CBL Signal transduction protein CBL CBL CBL2 RNFB55
Observed MW:	117 kDa
Uniprot ID :	P22681


Application

Reactivity: Human, Mouse


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:100

Image:



Immunohistochemistry of paraffin-embedded human prostate using FNab01317(CBL antibody) at dilution of 1:100



MCF7 cells were subjected to SDS PAGE followed by western blot with FNab01317(CBL antibody) at dilution of 1:1000