

CHD9 antibody

Product Information

Catalog No.:	FNab01645
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Acts as a transcriptional coactivator for PPARA and possibly other nuclear receptors. Proposed to be a ATP-dependent chromatin remodeling protein. Has DNA-dependent ATPase activity and binds to A/T-rich DNA. Associates with A/T-rich regulatory regions in promoters of genes that participate in the differentiation of progenitors during osteogenesis(By similarity).

Immunogen information


Immunogen:	chromodomain helicase DNA binding protein 9
Synonyms:	Chromodomain-helicase-DNA-binding protein 9 (CHD-9) ATP-dependent helicase CHD9 Chromatin-related mesenchymal modulator (CReMM) Chromatin-remodeling factor CHROM1 Kismet homolog 2 PPAR-alpha-interacting complex protein 320 kDa Peroxisomal proliferator-activated receptor A-interacting complex 320 kDa protein CHD9 KIAA0308 KISH2 PRIC320
Observed MW:	330 kDa
Uniprot ID :	Q3L8U1

Application


Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human kidney using FNab01645(CHD9 antibody) at dilution of 1:100



human brain tissue were subjected to SDS PAGE followed by western blot with FNab01645(CHD9 antibody) at dilution of 1:300