

COPS8 antibody

Product Information

Catalog No.:	FNab01874
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Component of the COP9 signalosome complex(CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin(Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IκappaBα/NFκBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

Immunogen information

Immunogen:	COP9 constitutive photomorphogenic homolog subunit 8(Arabidopsis)
Synonyms:	COP9 signalosome complex subunit 8 (SGN8, Signalosome subunit 8) COP9 homolog (hCOP9) JAB1-containing signalosome subunit 8 COPS8 CSN8
Observed MW:	23 kDa
Uniprot ID :	Q99627


Application

Reactivity:	Human, Mouse, Rat
-------------	-------------------


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:200-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded mouse brain tissue slide using FNab01874(COPS8 Antibody) at dilution of 1:50



mouse brain tissue were subjected to SDS PAGE followed by western blot with FNab01874(COPS8 Antibody) at dilution of 1:600