

DDB2 antibody

Product Information

Catalog No.:	FNab02286
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Required for DNA repair. Binds to DDB1 to form the UV-damaged DNA-binding protein complex(the UV-DDB complex). The UV-DDB complex may recognize UV-induced DNA damage and recruit proteins of the nucleotide excision repair pathway(the NER pathway) to initiate DNA repair. The UV-DDB complex preferentially binds to cyclobutane pyrimidine dimers(CPD), 6-4 photoproducts(6-4 PP), apurinic sites and short mismatches. Also appears to function as the substrate recognition module for the DCX(DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complex DDB1-CUL4-ROC1(also known as CUL4-DDB-ROC1 and CUL4-DDB-RBX1). The DDB1-CUL4-ROC1 complex may ubiquitinate histone H2A, histone H3 and histone H4 at sites of UV-induced DNA damage. The ubiquitination of histones may facilitate their removal from the nucleosome and promote subsequent DNA repair. The DDB1-CUL4-ROC1 complex also ubiquitinates XPC, which may enhance DNA-binding by XPC and promote NER. Isoform D1 and isoform D2 inhibit UV-damaged DNA repair.

Immunogen information

Immunogen:	damage-specific DNA binding protein 2, 48kDa
Synonyms:	DNA damage-binding protein 2 DDB p48 subunit (DDBb) Damage-specific DNA-binding protein 2 UV-damaged DNA-binding protein 2 (UV-DDB 2) DDB2
Observed MW:	55-60 kDa
Uniprot ID :	Q92466


Application

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IF

Recommended dilution: WB: 1:500-1:2000; IF: 1:20-1:200

Image:



HeLa cells were subjected to SDS PAGE followed by western blot with FNab02286(DDB2 Antibody) at dilution of 1:1000