

EIF3E antibody

Product Information

Catalog No.:	FNab02707
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Component of the eukaryotic translation initiation factor 3(eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAⁱ and eIF-5 to form the 43S preinitiation complex(43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Required for nonsense-mediated mRNA decay(NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins.

Immunogen information

Immunogen:	eukaryotic translation initiation factor 3, subunit E
Synonyms:	Eukaryotic translation initiation factor 3 subunit E (eIF3e) Eukaryotic translation initiation factor 3 subunit 6 Viral integration site protein INT-6 homolog eIF-3 p48 EIF3E EIF3S6 INT6
Observed MW:	48-52kd
Uniprot ID :	P60228


Application

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200; IF: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human pancreas cancer using FNab02707(EIF3E antibody) at dilution of 1:100



rat lymph tissue were subjected to SDS PAGE followed by western blot with FNab02707(EIF3E antibody) at dilution of 1:100