

FAAH antibody

Product Information

Catalog No.:	FNab02927
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background


Degrades bioactive fatty acid amides like oleamide, the endogenous cannabinoid, anandamide and myristic amide to their corresponding acids, thereby serving to terminate the signaling functions of these molecules. Hydrolyzes polyunsaturated substrate anandamide preferentially as compared to monounsaturated substrates.

Immunogen information


Immunogen:	fatty acid amide hydrolase
Synonyms:	Fatty-acid amide hydrolase 1 Anandamide amidohydrolase 1 Fatty acid ester hydrolase Oleamide hydrolase 1 FAAH FAAH1
Observed MW:	63 kDa
Uniprot ID :	O00519

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC
Recommended dilution:	WB: 1:500-1:1000; IHC: 1:20-1:200
Image:	



Immunohistochemistry of paraffin-embedded human testis tissue slide using FNab02927(FAAH Antibody) at dilution of 1:50



mouse liver tissue were subjected to SDS PAGE followed by western blot with FNab02927(FAAH Antibody) at dilution of 1:1000