

NR3C1 antibody

Product Information

| | |
|---------------|--|
| Catalog No.: | FNab03497 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background


None

Immunogen information


| | |
|--------------|--|
| Immunogen: | nuclear receptor subfamily 3, group C, member 1(glucocorticoid receptor) |
| Synonyms: | Glucocorticoid receptor (GR) Nuclear receptor subfamily 3 group C member 1 NR3C1 GRL |
| Observed MW: | 94-97 kDa |
| Uniprot ID : | P04150 |

Application


| | |
|-----------------------|---|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, WB,IHC, IF, IP |
| Recommended dilution: | WB: 1:500-1:2000; IHC: 1:20-1:200; IP: 1:200-1:1000; IF: 1:20-1:200 |
| Image: | |



Immunohistochemistry of paraffin-embedded human prostate cancer using FNab03497(GR antibody) at dilution of 1:100



IP Result of anti-Glucocorticoid receptor (IP:FNab03497, 4ug; Detection:FNab03497 1:500) with mouse heart tissue lysate 4400ug.



HeLa cells were subjected to SDS PAGE followed by western blot with FNab03497(Glucocorticoid receptor antibody) at dilution of 1:8000