

anti- HCK antibody

Product Information

Catalog No.:	FNab03785
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

The protein encoded by this gene is a member of the Src family of tyrosine kinases. This protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms with different subcellular distributions are produced due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) codon.

Immunogen information

Immunogen:	hemopoietic cell kinase
Synonyms:	None
Observed MW:	65 kDa
Uniprot ID :	P08631

Application

Reactivity:	Human, Mouse
Tested Application:	ELISA, WB, IHC, IF

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)


Tel : (0086)027-87384275

Fax: (0086)027-87800889


www.fn-test.com

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:10 - 1:100


Image:




Immunohistochemistry of paraffin-embedded human lung cancer using FNab03785(HCK Antibody) at dilution of 1:50



Immunofluorescence analysis of A549 cells using FNab03785(HCK antibody) at dilution of 1:50





Raji cells were subjected to SDS PAGE followed by western blot with FNab03785(HCK antibody) at dilution of 1:1000