

## HMOX1 antibody

### Product Information

Catalog No.:	FNab03937
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

### Background


Heme oxygenase cleaves the heme ring at the alpha methene bridge to form biliverdin. Biliverdin is subsequently converted to bilirubin by biliverdin reductase. Under physiological conditions, the activity of heme oxygenase is highest in the spleen, where senescent erythrocytes are sequestered and destroyed. Exhibits cytoprotective effects since excess of free heme sensitizes cells to undergo apoptosis.

### Immunogen information


Immunogen:	heme oxygenase(decycling) 1
Synonyms:	Heme oxygenase 1 (HO-1) Heme oxygenase 1 soluble form HMOX1 HO HO1
Observed MW:	33 kDa
Uniprot ID :	P09601

### Application


Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC, IP, IF
Recommended dilution:	WB: 1:500-1:1000; IP: 1:500-1:2000; IHC: 1:100-1:400; IF: 1:20-1:200
Image:	




Immunohistochemistry of paraffin-embedded human liver cancer tissue slide using FNab03937(HO-1 Antibody) at dilution of 1:200



Immunofluorescent analysis of ( -20°C Ethanol ) fixed HT-1080 cells using FNab03937 (HO-1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L)



IP Result of anti-HMOX1 (IP:FNab03937, 3ug; Detection:FNab03937 1:1000) with HeLa cells lysate 3000ug.



mouse liver tissue were subjected to SDS PAGE followed by western blot with FNab03937(HMOX1 antibody) at dilution of 1:600