

ICAM1 antibody

Product Information

Catalog No.: FNab04102

Size: 100µg
Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months(Avoid repeated freeze / thaw cycles.)

Background

ICAM-1(CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues(PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF- α , INF- γ , IL-1 and thrombin(PMID: 3086451; 9694714; 15979056).

Immunogen information

Immunogen: intercellular adhesion molecule 1

Synonyms: Intercellular adhesion molecule 1 (ICAM-1)|Major group rhinovirus

receptor|ICAM1

Observed MW: 85-95 kDa
Uniprot ID: P05362

Application


Reactivity: Mouse, Rat

Tested Application: ELISA, IF, WB, IHC, IP


Recommended dilution: WB: 1:500-1:1000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200




Image:



Immunohistochemistry of paraffin-embedded mouse lung using FNab04102(ICAM-1 antibody) at dilution of 1:50



IP Result of anti-ICAM-1 (IP:FNab04102, 4ug; Detection:FNab04102 1:1000) with Raji cells lysate 1320ug.



rat heart tissue were subjected to SDS PAGE followed by western blot with FNab04102(ICAM-1 antibody) at dilution of 1:800