

ICAM2 antibody

Product Information

Catalog No.:	FNab04105
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background


ICAM proteins are ligands for the leukocyte adhesion protein LFA-1(integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.

Immunogen information


Immunogen:	intercellular adhesion molecule 2
Synonyms:	Intercellular adhesion molecule 2 (ICAM-2) ICAM2
Observed MW:	55-80 kDa
Uniprot ID :	P13598

Application

Reactivity:	Human, Rat, Mouse
Tested Application:	ELISA, WB, IHC
Recommended dilution:	WB: 1:500-1:2000; IHC: 1:20-1:200
Image:	



Immunohistochemistry of paraffin-embedded human breast cancer using FNab04105(ICAM2 antibody) at dilution of 1:50



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab04105(ICAM2 antibody) at dilution of 1:500