

IL17RB antibody

Product Information

Catalog No.:	FNab04230
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

IL17RB, also named as CRL4, EVI27 and IL17BR, is a receptor for the proinflammatory cytokines IL17B and IL17E. IL17RB may play a role in controlling the growth and/or differentiation of hematopoietic cells. IL17RB has two isoforms with MW 56 kDa (isform 1) and 32 kDa (isform 2). The antibody is specific to IL17RB.


Immunogen information

Immunogen:	interleukin 17 receptor B
Synonyms:	Interleukin-17 receptor B (IL-17 receptor B, IL-17RB) Cytokine receptor-like 4 IL-17 receptor homolog 1 (IL-17Rh1, IL17Rh1) Interleukin-17B receptor (IL-17B receptor) IL17RB CRL4 EVI27 IL17BR
Observed MW:	33 kDa
Uniprot ID :	Q9NRM6


Application

Reactivity:	Human, Mouse
Tested Application:	ELISA, WB, IHC, IF, IP
Recommended dilution:	WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:50-1:500; IF: 1:10-1:100


Image:




Immunohistochemistry of paraffin-embedded human breast cancer using FNab04230(IL17RB antibody) at dilution of 1:100



Immunofluorescent analysis of HepG2 cells, using IL17RB antibody FNab04230(IL17RB antibody) at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP Result of anti-IL17RB (IP:FNab04230, 4ug; Detection:FNab04230 1:300) with PC-3 cells lysate 900ug.



human testis tissue were subjected to SDS PAGE followed by western blot with FNab04230(IL17RB antibody) at dilution of 1:500