

KIDINS220 antibody

Product Information

Catalog No.:	FNab04549
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Promotes a prolonged MAP-kinase signaling by neurotrophins through activation of a Rap1-dependent mechanism. Provides a docking site for the CRKL-C3G complex, resulting in Rap1-dependent sustained ERK activation. May play an important role in regulating postsynaptic signal transduction through the syntrophin-mediated localization of receptor tyrosine kinases such as EPHA4. In cooperation with SNTA1 can enhance EPHA4-induced JAK/STAT activation. Plays a role in nerve growth factor(NGF)-induced recruitment of RAPGEF2 to late endosomes and neurite outgrowth. May play a role in neurotrophin-and ephrin-mediated neuronal outgrowth and in axon guidance during neural development and in neuronal regeneration(By similarity). Modulates stress-induced apoptosis of melanoma cells via regulation of the MEK/ERK signaling pathway.

Immunogen information

Immunogen:	kinase D-interacting substrate, 220kDa
Synonyms:	Kinase D-interacting substrate of 220 kDa Ankyrin repeat-rich membrane-spanning protein KIDINS220 ARMS KIAA1250
Observed MW:	186-197 kDa
Uniprot ID :	Q9ULH0


Application

Reactivity:	Human, Mouse, Rat
-------------	-------------------

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



HeLa cells were subjected to SDS PAGE followed by western blot with FNab04549(KIDINS220 antibody) at dilution of 1:1000