

LCAT antibody

Product Information

Catalog No.:	FNab04722
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Central enzyme in the extracellular metabolism of plasma lipoproteins. Synthesized mainly in the liver and secreted into plasma where it converts cholesterol and phosphatidylcholines(lecithins) to cholesteryl esters and lysophosphatidylcholines on the surface of high and low density lipoproteins(HDLs and LDLs)(PubMed:10329423, PubMed:19065001, PubMed:26195816). The cholesterol ester is then transported back to the liver. Has a preference for plasma 16:0-18:2 or 18:0-18:2 phosphatidylcholines(PubMed:8820107). Also produced in the brain by primary astrocytes, and esterifies free cholesterol on nascent APOE-containing lipoproteins secreted from glia and influences cerebral spinal fluid(CSF) APOE-and APOA1 levels. Together with APOE and the cholesterol transporter ABCA1, plays a key role in the maturation of glial-derived, nascent lipoproteins. Required for remodeling high-density lipoprotein particles into their spherical forms(PubMed:10722751).

Immunogen information

Immunogen:	lecithin-cholesterol acyltransferase
Synonyms:	Phosphatidylcholine-sterol acyltransferase 1-alkyl-2-acetyl-glycerophosphocholine esterase Lecithin-cholesterol acyltransferase Phospholipid-cholesterol acyltransferase Platelet-activating factor acetylhydrolase (PAF acetylhydrolase) LCAT
Observed MW:	55 kDa
Uniprot ID :	P04180


Application

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



mouse testis tissue were subjected to SDS PAGE followed by western blot with FNab04722(LCAT antibody) at dilution of 1:500