

MED10 antibody

Product Information

Catalog No.:	FNab05086
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.


Immunogen information

Immunogen:	mediator complex subunit 10
Synonyms:	Mediator of RNA polymerase II transcription subunit 10 Mediator complex subunit 10 Transformation-related gene 17 protein (TRG-17) Transformation-related gene 20 protein (TRG-20) MED10
Observed MW:	16 kDa
Uniprot ID :	Q9BTT4

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB
Recommended dilution:	WB: 1:200 - 1:1000

Image:



COLO320 cells were subjected to SDS PAGE followed by western blot with FNab05086(MED10 Antibody) at dilution of 1:500