

## MED17 antibody

### Product Information

Catalog No.:	FNab05090
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

### Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

### Immunogen information

Immunogen:	mediator complex subunit 17
Synonyms:	Mediator of RNA polymerase II transcription subunit 17 Activator-recruited cofactor 77 kDa component (ARC77) Cofactor required for Sp1 transcriptional activation subunit 6 (CRSP complex subunit 6) Mediator complex subunit 17 Thyroid hormone receptor-associated protein complex 80 kDa component (Trap80) Transcriptional coactivator CRSP77 Vitamin D3 receptor-interacting protein complex 80 kDa component (DRIP80) MED17 ARC77 CRSP6 DRIP77 DRIP80 TRAP80
Observed MW:	80 kDa
Uniprot ID :	Q9NVC6

---


**Application**

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



human liver tissue were subjected to SDS PAGE followed by western blot with FNab05090(MED17 antibody) at dilution of 1:600