

MED26 antibody

Product Information

Catalog No.:	FNab05093
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional pre-initiation complex with RNA polymerase II and the general transcription factors.

Immunogen information

Immunogen:	mediator complex subunit 26
Synonyms:	Mediator of RNA polymerase II transcription subunit 26 Activator-recruited cofactor 70 kDa component (ARC70) Cofactor required for Sp1 transcriptional activation subunit 7 (CRSP complex subunit 7) Mediator complex subunit 26 Transcriptional coactivator CRSP70 MED26 ARC70 CRSP7
Observed MW:	66-70 kDa
Uniprot ID :	O95402


Application

Reactivity:	Human, Mouse, Rat
-------------	-------------------

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



mouse ovary tissue were subjected to SDS PAGE followed by western blot with FNab05093(MED26 antibody) at dilution of 1:500