

OCIAD1 antibody

Product Information

| | |
|---------------|--|
| Catalog No.: | FNab05958 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background


None

Immunogen information


| | |
|--------------|---|
| Immunogen: | OCIA domain containing 1 |
| Synonyms: | OCIA domain-containing protein 1 Ovarian cancer immunoreactive antigen domain containing 1 Ovarian carcinoma immunoreactive antigen OCIAD1 ASRIJ OCIA |
| Observed MW: | 29-35 kDa |
| Uniprot ID : | Q9NX40 |

Application


| | |
|-----------------------|---|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, IHC, IF, IP, WB |
| Recommended dilution: | WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200 |
| Image: | |



Immunohistochemistry of paraffin-embedded human liver using FNab05958(OCIAD1 antibody) at dilution of 1:100



IP Result of anti-OCIAD1 (IP:FNab05958, 4ug; Detection:FNab05958 1:300) with HeLa cells lysate 3200ug.



MCF7 cells were subjected to SDS PAGE followed by western blot with FNab05958(OCIAD1 antibody) at dilution of 1:300