

PMS2 antibody

Product Information

Catalog No.:	FNab06579
Size:	100µg
Form:	liquid
Purification:	Protein A+G purification
Purity:	\geq 95% as determined by SDS-PAGE
Host:	Mouse
Clonality:	monoclonal
Clone ID:	4B8
IsoType:	IgG1
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Component of the post-replicative DNA mismatch repair system(MMR). Heterodimerizes with MLH1 to form MutL alpha. DNA repair is initiated by MutS alpha(MSH2-MSH6) or MutS beta(MSH2-MSH6) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS-heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha(MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages.


Immunogen information

Immunogen:	PMS2 postmeiotic segregation increased 2
Synonyms:	Mismatch repair endonuclease PMS2 DNA mismatch repair protein PMS2 PMS1 protein homolog 2 PMS2 PMSL2
Observed MW:	100 kDa
Uniprot ID :	P54278




Application

Reactivity:Human, Mouse, RatTested Application:ELISA, WB, IHCRecommended dilution:WB: 1:500-1:2000; IHC: 1:50-1:500Image:



Immunohistochemistry of paraffin-embedded human cervical cancer tissue slide using FNab06579(PMS2 Antibody) at dilution of 1:400



A431 cells were subjected to SDS PAGE followed by western blot with FNab06579(PMS2 antibody) at dilution of 1:1000