

PSMD13 antibody

Product Information

Catalog No.:	FNab06885
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. Two transcripts encoding different isoforms have been described.

Immunogen information

Immunogen:	proteasome (prosome, macropain) 26S subunit, non-ATPase, 13
Synonyms:	26S proteasome non-ATPase regulatory subunit 13 26S proteasome regulatory subunit RPN9 26S proteasome regulatory subunit S11 26S proteasome regulatory subunit p40.5 PSMD13
Observed MW:	45 kDa
Uniprot ID :	Q9UNM6


Application

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:200

Image:



Immunohistochemistry of paraffin-embedded rat brain using FNab06885(PSMD13 antibody) at dilution of 1:100



mouse liver tissue were subjected to SDS PAGE followed by western blot with FNab06885(PSMD13 antibody) at dilution of 1:1000