

PSMD4 antibody

Product Information

| | |
|---------------|--|
| Catalog No.: | FNab06890 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Protein A+G purification |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Mouse |
| Clonality: | monoclonal |
| Clone ID: | 0A11 |
| IsoType: | IgG1 |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background


Binds and presumably selects ubiquitin-conjugates for destruction. Displays selectivity for longer polyubiquitin chains. Modulates intestinal fluid secretion.

Immunogen information


| | |
|--------------|---|
| Immunogen: | proteasome(prosome, macropain) 26S subunit, non-ATPase, 4 |
| Synonyms: | 26S proteasome non-ATPase regulatory subunit 4 26S proteasome regulatory subunit RPN10 26S proteasome regulatory subunit S5A Antisecretory factor 1 (AF, ASF) Multiubiquitin chain-binding protein PSMD4 MCB1 |
| Observed MW: | 50 kDa |
| Uniprot ID : | P55036 |

Application

| | |
|-----------------------|---|
| Reactivity: | Human |
| Tested Application: | ELISA, WB, IF, IHC |
| Recommended dilution: | WB: 1:500-1:2000; IF: 1:20-1:200; IHC: 1:20-1:200 |
| Image: | |



Immunohistochemistry of paraffin-embedded human testis tissue slide using FNab06890(PSMD4 Antibody) at dilution of 1:200



HeLa cells were subjected to SDS PAGE followed by western blot with FNab06890(PSMD4 Antibody) at dilution of 1:1000