

RBM17 antibody

Product Information

| | |
|---------------|--|
| Catalog No.: | FNab07166 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background


Splice factor that binds to the single-stranded 3'AG at the exon/intron border and promotes its utilization in the second catalytic step. Involved in the regulation of alternative splicing and the utilization of cryptic splice sites. Promotes the utilization of a cryptic splice site created by the beta-110 mutation in the HBB gene. The resulting frameshift leads to sickle cell anemia.

Immunogen information

| | |
|--------------|--|
| Immunogen: | RNA binding motif protein 17 |
| Synonyms: | Splicing factor 45 45 kDa-splicing factor RNA-binding motif protein 17 RBM17 SPF45 |
| Observed MW: | 45 kDa |
| Uniprot ID : | Q96I25 |

Application

| | |
|-----------------------|----------------------------------|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, WB, IF |
| Recommended dilution: | WB: 1:500-1:2000; IF: 1:20-1:200 |
| Image: | |



HeLa cells were subjected to SDS PAGE followed by western blot with FNab07166(RBM17 antibody) at dilution of 1:3000