

RBM17 antibody

Product Information

Catalog No.:	FNab07167
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background


Splice factor that binds to the single-stranded 3'AG at the exon/intron border and promotes its utilization in the second catalytic step. Involved in the regulation of alternative splicing and the utilization of cryptic splice sites. Promotes the utilization of a cryptic splice site created by the beta-110 mutation in the HBB gene. The resulting frameshift leads to sickle cell anemia.

Immunogen information

Immunogen:	RNA binding motif protein 17
Synonyms:	Splicing factor 45 45 kDa-splicing factor RNA-binding motif protein 17 RBM17 SPF45
Observed MW:	45 kDa
Uniprot ID :	Q96I25

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB
Recommended dilution:	WB: 1:500-1:2000
Image:	



HeLa cells were subjected to SDS PAGE followed by western blot with FNab07167(RBM17 antibody) at dilution of 1:3000