

RBPJ antibody

Product Information

Catalog No.:	FNab07196
Size:	100µg
Form:	liquid
Purification:	Protein A+G purification
Purity:	≥95% as determined by SDS-PAGE
Host:	Mouse
Clonality:	monoclonal
Clone ID:	2A10
IsoType:	IgG1
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background

Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins(Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA. Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions(4% oxygen)(PubMed:23303788).

Immunogen information

Immunogen:	recombination signal binding protein for immunoglobulin kappa J region
Synonyms:	Recombining binding protein suppressor of hairless CBF-1 J kappa-recombination signal-binding protein RBP-J kappa (RBP-J, RBP-JK) Renal carcinoma antigen NY-REN-30 RBPJ IGKJRB IGKJRB1 RBPJK RBPSUH
Observed MW:	60 kDa
Uniprot ID :	Q06330


Application

Reactivity: Human

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab07196(RBPJ antibody) at dilution of 1:1000