

RPA1 antibody

Product Information

Catalog No.:	FNab07391	
Size:	100µg	
Form:	liquid	
Purification:	Immunogen affinity purified	
Purity:	\geq 95% as determined by SDS-PAGE	
Host:	Rabbit	
Clonality:	polyclonal	
Clone ID:	None	
IsoType:	IgG	
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)	

Background

As part of the heterotrimeric replication protein A complex(RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage(PubMed:9430682). In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response(PubMed:24332808). It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage(PubMed:17765923). Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair(PubMed:7697716). Plays also a role in base excision repair(BER) probably through interaction with UNG(PubMed:9765279). Through RFWD3 may activate CHEK1 and play a role in replication checkpoint control. Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance(PubMed:17959650). As part of the alternative replication protein A complex, aRPA, binds single-stranded DNA and probably plays a role in DNA repair. Compared to the RPA2-containing, canonical RPA complex, may not support chromosomal DNA replication and cell cycle progression through S-phase. The aRPA may not promote efficient priming by DNA polymerase alpha but could support DNA synthesis by polymerase delta in presence of PCNA and replication factor C(RFC), the dual incision/excision reaction of nucleotide excision repair and RAD51-dependent strand exchange(PubMed:19996105).



Immunogen information

Immunogen:	replication protein A1, 70kDa
Synonyms:	Replication protein A 70 kDa DNA-binding subunit (RP-A p70) Replication factor A protein 1 (RF-A protein 1) Single-stranded DNA-binding protein Replication protein A 70 kDa DNA-binding subunit, N-terminally processed RPA1 REPA1 RPA70
Observed MW:	70 kDa
Uniprot ID :	P27694

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB
Recommended dilution	:WB: 1:500 - 1:2000
Image:	



Hela cells were subjected to SDS PAGE followed by western blot with FNab07391(RPA1 antibody) at dilution of 1:1000