

## SSX2IP antibody

### Product Information

Catalog No.:	FNab08258
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

### Background

Belongs to an adhesion system, which plays a role in the organization of homotypic, interneuronal and heterotypic cell-cell adherens junctions(AJs). May connect the nectin-afadin and E-cadherin-catenin system through alpha-actinin and may be involved in organization of the actin cytoskeleton at AJs through afadin and alpha-actinin(By similarity). Involved in cell movement: localizes at the leading edge of moving cells in response to PDGF and is required for the formation of the leading edge and the promotion of cell movement, possibly via activation of Rac signaling(By similarity). Acts as a centrosome maturation factor, probably by maintaining the integrity of the pericentriolar material and proper microtubule nucleation at mitotic spindle poles. The function seems to implicate at least in part WRAP73; the SSX2IP:WRAP73 complex is proposed to act as regulator of spindle anchoring at the mitotic centrosome(PubMed:23816619, PubMed:26545777). Involved in ciliogenesis(PubMed:24356449). It is required for targeted recruitment of the BBSome, CEP290, RAB8, and SSTR3 to the cilia(PubMed:24356449).

### Immunogen information

Immunogen:	synovial sarcoma, X breakpoint 2 interacting protein
Synonyms:	Afadin- and alpha-actinin-binding protein (ADIP) Afadin DIL domain-interacting protein SSX2-interacting protein SSX2IP KIAA0923
Observed MW:	64 kDa
Uniprot ID :	Q9Y2D8

---


**Application**

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



HEK-293 cells were subjected to SDS PAGE followed by western blot with FNab08258(SSX2IP antibody) at dilution of 1:500