

STAMBP antibody

Product Information

Catalog No.: FNab08285

Size: 100µg
Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Cytokine-mediated signal transduction in the JAK-STAT cascade requires the involvement of adaptor molecules. One such signal-transducing adaptor molecule contains an SH3 domain that is required for induction of MYC and cell growth. The protein encoded by this gene binds to the SH3 domain of the signal-transducing adaptor molecule, and plays a critical role in cytokine-mediated signaling for MYC induction and cell cycle progression. Multiple alternatively spliced transcript variants encoding the same protein isoform have been found for this gene.

Immunogen information

Immunogen: STAM binding protein

Synonyms: STAM-binding protein Associated molecule with the SH3 domain of

STAM|Endosome-associated ubiquitin isopeptidase|STAMBP|AMSH

Observed MW: 48 kDa
Uniprot ID: 095630


Application

Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC, IF


Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:100



Image:



Immunohistochemistry of paraffin-embedded human prostate cancer using FNab08285(STAMBP antibody) at dilution of 1:100



mouse brain tissue were subjected to SDS PAGE followed by western blot with FNab08285(STAMBP antibody) at dilution of 1:1000