

STARD10 antibody

Product Information

Catalog No.:	FNab08289
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background


May play metabolic roles in sperm maturation or fertilization(By similarity). Phospholipid transfer protein that preferentially selects lipid species containing a palmitoyl or stearoyl chain on the sn-1 and an unsaturated fatty acyl chain(18:1 or 18:2) on the sn-2 position. Able to transfer phosphatidylcholine(PC) and phosphatidylethanolamine(PE) between membranes.

Immunogen information


Immunogen:	StAR-related lipid transfer(START) domain containing 10
Synonyms:	START domain-containing protein 10 (StARD10) Antigen NY-CO-28 PCTP-like protein (PCTP-L) Serologically defined colon cancer antigen 28 StAR-related lipid transfer protein 10 STARD10 SDCCAG28
Observed MW:	35 kDa
Uniprot ID :	Q9Y365

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC
Recommended dilution:	WB: 1:500-1:2000; IHC: 1:20-1:200
Image:	



Immunohistochemistry of paraffin-embedded human liver using FNab08289(STARD10 antibody) at dilution of 1:100



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab08289(STARD10 antibody) at dilution of 1:1500