

STARD8 antibody

Product Information

Catalog No.:	FNab08293
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.)

Background


Accelerates GTPase activity of RHOA and CDC42, but not RAC1. Stimulates the hydrolysis of phosphatidylinositol 4,5-bisphosphate by PLCD1.

Immunogen information


Immunogen:	StAR-related lipid transfer(START) domain containing 8
Synonyms:	StAR-related lipid transfer protein 8 Deleted in liver cancer 3 protein (DLC-3) START domain-containing protein 8 (StARD8) START-GAP3 STARD8 DLC3 KIAA0189
Observed MW:	115-120 kDa
Uniprot ID :	Q92502

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC
Recommended dilution:	WB: 1:500-1:2000; IHC: 1:20-1:200
Image:	



Immunohistochemistry of paraffin-embedded human stomach cancer tissue slide using FNab08293(STARD8 Antibody) at dilution of 1:50



HEK-293 cells were subjected to SDS PAGE followed by western blot with FNab08293(STARD8 antibody) at dilution of 1:400