

anti- STAT5B antibody

Product Information

Catalog No.: FNab08305

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Carries out a dual function: signal transduction and activation of transcription. Mediates cellular responses to the cytokine KITLG/SCF and other growth factors. Binds to the GAS element and activates PRL-induced transcription.

Immunogen information

Immunogen: signal transducer and activator of transcription 5B

Synonyms: None
Observed MW: 90 kDa
UniprotID: P51692

Application

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IHC, IF, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200

Image:


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>

1






Immunohistochemistry of paraffin-embedded human lung cancer using FNab08305(STAT5B antibody) at dilution of 1:100



IP Result of anti-STAT5B (IP:FNab08305, 4ug; Detection:FNab08305 1:500) with HeLa cells lysate 1200ug.





HeLa cells were subjected to SDS PAGE followed by western blot with FNab08305(STAT5B antibody) at dilution of 1:1000

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com