

STAU1 antibody

Product Information

Catalog No.:	FNab08312
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Staufen is a member of the family of double-stranded RNA (dsRNA)-binding proteins involved in the transport and/or localization of mRNAs to different subcellular compartments and/or organelles. These proteins are characterized by the presence of multiple dsRNA-binding domains which are required to bind RNAs having double-stranded secondary structures. The human homologue of staufen encoded by STAU, in addition contains a microtubule-binding domain similar to that of microtubule-associated protein 1B, and binds tubulin. The STAU gene product has been shown to be present in the cytoplasm in association with the rough endoplasmic reticulum (RER), implicating this protein in the transport of mRNA via the microtubule network to the RER, the site of translation. Five transcript variants resulting from alternative splicing of STAU gene and encoding three isoforms have been described. Three of these variants encode the same isoform, however, differ in their 5'UTR.

Immunogen information

Immunogen:	staufen, RNA binding protein, homolog 1 (Drosophila)
Synonyms:	Double-stranded RNA-binding protein Staufen homolog 1 STAU1 STAU
Observed MW:	58 kDa
Uniprot ID :	O95793


Application

Reactivity:	Human, Mouse, Rat
-------------	-------------------


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:100

Image:



Immunohistochemistry of paraffin-embedded human esophagus using FNab08312(STAU1 antibody) at dilution of 1:100



SW480 cells were subjected to SDS PAGE followed by western blot with FNab08312(STAU1 antibody) at dilution of 1:1000