

TMF1-Specific antibody

Product Information

| | |
|---------------|--|
| Catalog No.: | FNab08796 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background


None

Immunogen information


| | |
|--------------|--|
| Immunogen: | TATA element modulatory factor 1 |
| Synonyms: | TATA element modulatory factor (TMF) Androgen receptor coactivator 160 kDa protein Androgen receptor-associated protein of 160 kDa TMF1 ARA160 |
| Observed MW: | 150 kDa |
| Uniprot ID : | P82094 |

Application


| | |
|-----------------------|---|
| Reactivity: | Human |
| Tested Application: | ELISA, WB, IHC, IP |
| Recommended dilution: | WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200 |
| Image: | |



Immunohistochemistry of paraffin-embedded human cervical cancer using FNab08796(TMFI-Specific antibody) at dilution of 1:100



IP Result of anti-TMF1-Specific (IP:FNab08796, 4ug; Detection:FNab08796 1:300) with HepG2 cells lysate 3600ug.



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab08796(TMFI-Specific antibody) at dilution of 1:300