

TNFAIP8L2 antibody

Product Information

Catalog No.: FNab08820

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months(Avoid repeated freeze / thaw cycles.)

Background

Acts as a negative regulator of innate and adaptive immunity by maintaining immune homeostasis. Negative regulator of Toll-like receptor and T-cell receptor function. Prevents hyperresponsiveness of the immune system and maintains immune homeostasis. Inhibits JUN/AP1 and NF-kappa-B activation. Promotes Fas-induced apoptosis(By similarity).

Immunogen information

Immunogen: tumor necrosis factor, alpha-induced protein 8-like 2

Synonyms: Tumor necrosis factor alpha-induced protein 8-like protein 2 (TIPE2,

TNF alpha-induced protein 8-like protein 2, TNFAIP8-like protein

2)|Inflammation factor protein 20|TNFAIP8L2

Observed MW: 18-21 kDa Uniprot ID: Q6P589

Application


Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IHC, IP, IF


Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:2000; IHC: 1:20-1:200; IF: 1:20-1:200

Image:






Immunohistochemistry of paraffin-embedded human lymphoma using FNab08820(TNFAIP8L2 antibody) at dilution of 1:50



IP Result of anti-TIPE2 (IP:FNab08820, 3ug; Detection:FNab08820 1:500) with mouse cerebellum tissue lysate 4000ug.



mouse cerebellum tissue were subjected to SDS PAGE followed by western blot with FNab08820(TNFAIP8L2 antibody) at dilution of 1:500