

TTF2 antibody

Product Information

| | |
|---------------|--|
| Catalog No.: | FNab09092 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months(Avoid repeated freeze / thaw cycles.) |

Background


DsDNA-dependent ATPase which acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template. May contribute to mitotic transcription repression. May also be involved in pre-mRNA splicing.

Immunogen information


| | |
|--------------|---|
| Immunogen: | transcription termination factor, RNA polymerase II |
| Synonyms: | Transcription termination factor 2 Lodestar homolog RNA polymerase II termination factor Transcription release factor 2 (F2, HuF2) TTF2 |
| Observed MW: | 130 kDa |
| Uniprot ID : | Q9UNY4 |

Application


| | |
|-----------------------|---|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, WB, IHC, IF, IP |
| Recommended dilution: | WB: 1:500-1:5000; IP: 1:200-1:1000; IF: 1:20-1:200; IHC: 1:20-1:200 |
| Image: | |



Immunohistochemistry of paraffin-embedded human ovary tumor using FNab09092(TTF2 antibody) at dilution of 1:100



IP Result of anti-TTF2 (IP:FNab09092, 4ug; Detection:FNab09092 1:500) with A2780 cells lysate 840ug.



A2780 cells were subjected to SDS PAGE followed by western blot with FNab09092(TTF2 antibody) at dilution of 1:500