

UBE2T antibody

Product Information

Catalog No.: FNab09185

Size: 100µg
Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months(Avoid repeated freeze / thaw cycles.)

Background

The ubiquitin(Ub)-mediated protein degradation pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub to specific protein substrates. The first step requires ATP-dependent activation of the C-terminus of Ub and the assembly of multi-Ubs by Ub-activating enzyme E1. The ubiquitin-conjugating enzyme E2, catalytic(UBCc) domain, is then conjugated to Ubs, through a thiol-ester linkage between a conserved cysteine and the C-terminus of Ub, to generate an intermediate Ub-E2 complex. Then the E3, a ligase, catalyzes the transfer of Ub from E2 to the appropriate substrate. This pathway regulates many fundamental cellular processes. There are also other E2s which form thiol-ester linkages without the use of E3s as well as several UBC homologs(TSG101, Mms2, Croc-1 and similar proteins), which lack the active site cysteine essential for ubiquitination and appear to function in DNA repair pathways.

Immunogen information

Immunogen: ubiquitin-conjugating enzyme E2T(putative)

Synonyms: Ubiquitin-conjugating enzyme E2 T|Cell proliferation-inducing gene 50

protein|E2 ubiquitin-conjugating enzyme T|Ubiquitin carrier protein

T|Ubiquitin-protein ligase T|UBE2T

Observed MW: 24 kDa
Uniprot ID: Q9NPD8




Application


Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IP

Recommended dilution: WB: 1:200-1:2000; IP: 1:200-1:2000

Image:



IP Result of anti-UBE2T/HSPC150 (IP:FNab09185, 3ug; Detection:FNab09185 1:500) with HeLa cells lysate 3000ug.



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab09185(UBE2T antibody) at dilution of 1:800