

# anti- Beta Tubulin antibody

### **Product Information**

Catalog No.:	FNab09872
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3 , -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

#### Background

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha, beta, and gamma tubulins have all been used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.

#### **Immunogen information**

Immunogen:	tubulin, beta 3
Synonyms:	beta 4, beta tubulin, MC1R, TUBB, tubulin
Calculated MW:	50-55 kDa
Uniprot ID :	Q13509

## Application

Specificity:	Human, Mouse, Rat; other species are not tested. Please decide the specificity by homology	
Tested Application:	ELISA, WB, IHC	
Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200		


#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave.East Lake High-Tech Development Zone.Wuhan,<br/>Hubei, China(430206)Tel :( 0086)027-87384275Fax: (0086)027-87800889www.fn-test.com


1



## Image:



Immunohistochemistry of paraffin-embedded human liver cancer using FNab09872(beta Tubulin Antibody) at dilution of 1:50



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab09872(beta Tubulin antibody) at dilution of 1:1000

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)